agent(s) described under §610.40(a) or reactive with a serological test for syphilis under §610.40(i), may serve as an autologous donor under §610.40(d).

- (b) A deferred donor subsequently may be found to be suitable as a donor of blood or blood components by a requalification method or process found acceptable for such purposes by FDA. Such a donor is considered no longer deferred.
- (c) You must comply with the requirements under §§610.46 and 610.47 when a donor tests reactive by a screening test for HIV or HCV required under §610.40(a) and (b), or when you are aware of other reliable test results or information indicating evidence of HIV or HCV infection.

[66 FR 31164, June 11, 2001, as amended at 72 FR 48798, Aug. 24, 2007]

§ 610.42 Restrictions on use for further manufacture of medical devices.

- (a) In addition to labeling requirements in subchapter H of this chapter, when a medical device contains human blood or a blood component as a component of the final device, and the human blood or blood component was found to be reactive by a screening test performed under §610.40(a) and (b) or reactive for syphilis under §610.40(i). then you must include in the device labeling a statement of warning indicating that the product was manufactured from a donation found to be reactive by a screening test for evidence of infection due to the identified communicable disease agent(s).
- (b) FDA may approve an exception or alternative to the statement of warning required in paragraph (a) of this section based on evidence that the reactivity of the human blood or blood component in the medical device presents no significant health risk through use of the medical device.

[66 FR 31164, June 11, 2001]

§ 610.44 Use of reference panels by manufacturers of test kits.

(a) When available and appropriate to verify acceptable sensitivity and specificity, you, a manufacturer of test kits, must use a reference panel you obtain from FDA or from an FDA designated source to test lots of the following

products. You must test each lot of the following products, unless FDA informs you that less frequent testing is appropriate, based on your consistent prior production of products of acceptable sensitivity and specificity:

- (1) A test kit approved for use in testing donations of human blood and blood components for evidence of infection due to communicable disease agents listed in §610.40(a); and
- (2) Human immunodeficiency virus (HIV) test kit approved for use in the diagnosis, prognosis, or monitoring of this communicable disease agent.
- (b) You must not distribute a lot that is found to be not acceptable for sensitivity and specificity under §610.44(a). FDA may approve an exception or alternative to this requirement. Applicants must submit such requests in writing. However, in limited circumstances, such requests may be made orally and permission may be given orally by FDA. Oral requests and approvals must be promptly followed by written requests and written approvals.

[66 FR 31164, June 11, 2001]

§ 610.46 Human immunodeficiency virus (HIV) "lookback" requirements.

- (a) If you are an establishment that collects Whole Blood or blood components, including Source Plasma and Source Leukocytes, you must establish, maintain, and follow an appropriate system for the following actions:
- (1) Within 3 calendar days after a donor tests reactive for evidence of human immunodeficiency virus (HIV) infection when tested under §610.40(a) and (b) or when you are made aware of other reliable test results or information indicating evidence of HIV infection, you must review all records required under §606.160(d) of this chapter, to identify blood and blood components previously donated by such a donor. For those identified blood and blood components collected:
- (i) Twelve months and less before the donor's most recent nonreactive screening tests, or
- (ii) Twelve months and less before the donor's reactive direct viral detection test, e.g., nucleic acid test or HIV

§610.46

p24 antigen test, and nonreactive antibody screening test, whichever is the lesser period, you must:

- (A) Quarantine all previously collected in-date blood and blood components identified under paragraph (a)(1) of this section if intended for use in another person or for further manufacture into injectable products, except pooled blood components intended solely for further manufacturing into products that are manufactured using validated viral clearance procedures; and
- (B) Notify consignees to quarantine all previously collected in-date blood and blood components identified under paragraph (a)(1) of this section if intended for use in another person or for further manufacture into injectable products, except pooled blood components intended solely for further manufacturing into products that are manufactured using validated viral clearance procedures;
- (2) You must perform a supplemental (additional, more specific) test for HIV as required under §610.40(e) of this chapter on the reactive donation.
- (3) You must notify consignees of the supplemental (additional, more specific) test results for HIV, or the results of the reactive screening test if there is no available supplemental test that is approved for such use by FDA, or if under an investigational new drug application (IND) or investigational device exemption (IDE), is exempted for such use by FDA, within 45 calendar days after the donor tests reactive for evidence of HIV infection under §610.40(a) and (b) of this chapter. Notification of consignees must include the test results for blood and blood components identified under paragraph (a)(1) of this section that were previously collected from donors who later test reactive for evidence of HIV infection.
- (4) You must release from quarantine, destroy, or relabel quarantined in-date blood and blood components, consistent with the results of the supplemental (additional, more specific) test performed under paragraph (a)(2) of this section or the results of the reactive screening test if there is no available supplemental test that is approved for such use by FDA, or if under an IND or IDE, exempted for such use by FDA.

- (b) If you are a consignee of Whole Blood or blood components, including Source Plasma and Source Leukocytes, you must establish, maintain, and follow an appropriate system for the following actions:
- (1) You must quarantine all previously collected in-date blood and blood components identified under paragraph (a)(1) of this section, except pooled blood components intended solely for further manufacturing into products that are manufactured using validated viral clearance procedures, when notified by the collecting establishment.
- (2) You must release from quarantine, destroy, or relabel quarantined in-date blood and blood components consistent with the results of the supplemental (additional, more specific) test performed under paragraph (a)(2) of this section, or the results of the reactive screening test if there is no available supplemental test that is approved for such use by FDA, or if under an IND or IDE, is exempted for such use by FDA.
- (3) When the supplemental (additional, more specific) test for HIV is positive or when the screening test is reactive and there is no available supplemental test that is approved for such use by FDA, or if under an IND or IDE is exempted for such use by FDA, you must notify transfusion recipients of previous collections of blood and blood components at increased risk of transmitting HIV infection, or the recipient's physician of record, of the need for recipient HIV testing and counseling. You must notify the recipient's physician of record or a legal representative or relative if the recipient is a minor, deceased, adjudged incompetent by a State court, or, if the recipient is competent but State law permits a legal representative or relative to receive information on behalf of the recipient. You must make reasonable attempts to perform the notification within 12 weeks after receiving the supplemental (additional, more specific) test results for evidence of HIV infection from the collecting establishment, or after receiving the donor's reactive screening test result for HIV if there is no available supplemental test that is approved for such use by FDA, or if

under an IND or IDE is exempted for such use by FDA.

(c) Actions under this section do not constitute a recall as defined in §7.3 of this chapter.

[72 FR 48799, Aug. 24, 2007]

§610.47 Hepatitis C virus (HCV) "lookback" requirements.

- (a) If you are an establishment that collects Whole Blood or blood components, including Source Plasma and Source Leukocytes, you must establish, maintain, and follow an appropriate system for the following actions:
- (1) Within 3 calendar days after a donor tests reactive for evidence of hepatitis C virus (HCV) infection when tested under §610.40(a) and (b) of this chapter or when you are made aware of other reliable test results or information indicating evidence of HCV infection, you must review all records required under §606.160(d) of this chapter, to identify blood and blood components previously donated by such a donor. For those identified blood and blood components collected:
- (i) Twelve months and less before the donor's most recent nonreactive screening tests, or
- (ii) Twelve months and less before the donor's reactive direct viral detection test, e.g., nucleic acid test and nonreactive antibody screening test, whichever is the lesser period, you must:
- (A) Quarantine all previously collected in-date blood and blood components identified under paragraph (a)(1) of this section if intended for use in another person or for further manufacture into injectable products, except pooled blood components intended solely for further manufacturing into products that are manufactured using validated viral clearance procedures; and
- (B) Notify consignees to quarantine all previously collected in-date blood and blood components identified under paragraph (a)(1) of this section if intended for use in another person or for further manufacture into injectable products, except pooled blood components intended solely for further manufacturing into products that are manufactured using validated viral clearance procedures;

- (2) You must perform a supplemental (additional, more specific) test for HCV as required under §610.40(e) on the reactive donation.
- (3) You must notify consignees of the supplemental (additional, more specific) test results for HCV, or the results of the reactive screening test if there is no available supplemental test that is approved for such use by FDA, or if under an investigational new drug application (IND) or investigational device exemption (IDE), is exempted for such use by FDA, within 45 calendar days after the donor tests reactive for evidence of HCV infection under §610.40(a) and (b). Notification of consignees must include the test results for blood and blood components identified under paragraph (a)(1) of this section that were previously collected from donors who later test reactive for evidence of HCV infection.
- (4) You must release from quarantine, destroy, or relabel quarantined in-date blood and blood components consistent with the results of the supplemental (additional, more specific) test performed under paragraph (a)(2) of this section, or the results of the reactive screening test if there is no available supplemental test that is approved for such use by FDA, or if under an IND or IDE, exempted for such use by FDA.
- (b) If you are a consignee of Whole Blood or blood components, including Source Plasma or Source Leukocytes, you must establish, maintain, and follow an appropriate system for the following actions:
- (1) You must quarantine all previously collected in-date blood and blood components identified under paragraph (a)(1) of this section, except pooled blood components intended solely for further manufacturing into products that are manufactured using validated viral clearance procedures, when notified by the collecting establishment.
- (2) You must release from quarantine, destroy, or relabel quarantined in-date blood and blood components, consistent with the results of the supplemental (additional, more specific test performed under paragraph (a)(2) of this section, or the results of the reactive screening test if there is no